

EDGE SUM LABELING ON PENDENT GRAPHS

S. Alice Pappa, R.Sweetly, *G.J. Jeba Selvi Kavitha

¹Department of Mathematics, Nazareth Margoschis College at Pillaiyanmanai Tuticorin-Affiliated to Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli – 627012. Tamil Nadu, India
Email alicestephen8979@gmail.com

²Department of Mathematics, Nazareth Margoschis College at Pillaiyanmanai Tuticorin-Affiliated to Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli – 627012. Tamil Nadu India.
Email sweetydaniel65@gmail.com

³Department of Mathematics, St. Joseph's College of Arts and Science for Women-Hours and Research Scholar, Reg. No. 19232142092007, Department of Mathematics, Nazareth Margoschis College At Pillaiyanmanai Tuticorin-Affiliated to Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli – 627012, Tamil Nadu, India
Email jebaselvikavitha@gmail.com

Abstract

Let $G(V, E)$ be a graph with p vertices and q edges. Let $f: V(G) \rightarrow \{0, 1, 2, \dots, p-1\}$ be a bijection such that the induced function $f^*: E(G) \rightarrow N$ defined by $f_{sum}^*(uv) = f(u) + f(v)$ for every $uv \in E(G)$. If f_{sum}^* is injective then f_{sum}^* is called an edge sum graph labeling of G . A graph which admits sum graph labeling is called edge sum graph.

1. Introduction

Every graph in this paper are simple finite, undirected and non-trivial graph $G(V, E)$ with vertex set V and Edge set E . For graph theoretic terminology we refer to Harary. R. Ponraj, J. Vijaya Xavier Parthipan and R. Kala defined pair sum labeling. We introduce edge sum labeling on pendent graphs

Definition 2.1

Pendent graph with a vertex is obtained by appending a pendent edges of each vertex of a path.

Definition 2.2

Pendent graph with 2 vertex is obtained by appending a pendent edges of each vertex of a path.

Definition 2.3

Pendent graph with 3 vertex is obtained by appending a pendent edges of each vertex of a path.

Theorem 3.1

Pendent graph with a pendent is an edge sum graph.

Proof:

Let $V = \{v_i / 1 \leq i \leq n\} \cup \{u_i / 1 \leq i \leq n\}$ be the vertex set of simple pendant graph with 1 pendant and $E = \bigcup_{i=1}^n E_i$

Let $E_1 = \{v_i v_{i+1} / 1 \leq i \leq n-1\}$

$E_2 = \{v_i u_i / 1 \leq i \leq n\}$

Here $|V(G)|=2n$

$|E(G)|=2n-1$

Define $f:V \rightarrow \{0, 1, 2, \dots, p-1\}$ by

$$f(v_i) = \{2(i-1)/1 \leq i \leq n\}$$

$$f(u_i) = \{2i-1/1 \leq i \leq n\}$$

Then the corresponding edge label are as follows

$$f_{sum}^*(v_i v_{i+1}) = f(v_i) + f(v_{i+1})$$

$$= 2(i-1) + 2(i+1-1)$$

$$= 2i - 2 + 2i$$

$$= 4i - 2$$

$$f_{sum}^*(v_i u_i) = f(v_i) + f(u_i)$$


$$= 2(i-1) + 2i - 1$$

$$= 2i - 2 + 2i - 1$$

$$= 4i - 3$$

Clearly the edge labels are distinct.

Therefore, pendant graph with a pendant is a edge sum graph.



Theorem 3.2

Pendant graph with 2 pendant is a edge sum graph.

Proof

Let $V = \{v_i \mid 1 \leq i \leq n\} \cup \{u_i \mid 1 \leq i \leq n\} \cup \{w_i \mid 1 \leq i \leq n\}$

be the vertex set of simple pendant graph with 2 pendant and let

$$E = \bigcup_{i=1}^3 E_i$$

$$E_1 = \{v_i v_{i+1} \mid 1 \leq i \leq n-1\}$$

$$E_2 = \{v_i u_i \mid 1 \leq i \leq n\}$$

$$E_3 = \{v_i w_i | 1 \leq i \leq n\}$$

Here $|V(G)| = 3n$ and $|E(G)| = 3n-1$

Define $f: V \rightarrow \{0, 1, 2, \dots, p-1\}$ by

$$f(v_i) = \{3(i-1) | 1 \leq i \leq n\}$$

$$f(u_i) = \{3i-2 | 1 \leq i \leq n\}$$

$$f(w_i) = \{3i-1 | 1 \leq i \leq n\}$$

Then the corresponding edge labeling is as follows


$$\begin{aligned} f_{sum}^*(v_i v_{i+1}) &= f(v_i) + f(v_{i+1}) \\ &= 3(i-1) + 3(i+1-1) \\ &= 3(i-1) + 3i \\ &= 6i-3 \end{aligned}$$

$$\begin{aligned} f_{sum}^*(v_i u_i) &= f(v_i) + f(u_i) \\ &= 3(i-1) + 3i-2 \\ &= 3i-3 + 3i-2 \\ &= 6i-5 \end{aligned}$$

$$\begin{aligned} f_{sum}^*(v_i w_i) &= f(v_i) + f(w_i) \\ &= 3(i-1) + 3i-1 \\ &= 3i-3 + 3i-1 \\ &= 6i-4 \end{aligned}$$

Clearly the edge labels are distinct.

Therefore pendant graph with 2 pendant is a edge sum graph.



Theorem 3.3

Pendant graph with 3 pendant is a edge sum graph.

Proof:

$$\text{Let } V = \{v_i \mid 1 \leq i \leq n\} \cup \{u_i \mid 1 \leq i \leq n\} \cup \{w_i \mid 1 \leq i \leq n\} \cup \{x_i \mid 1 \leq i \leq n\}$$

be the vertex set of pendant graph with 3 pendant and

$$E = \bigcup_{i=1}^4 E_i$$

$$\text{Let } E_1 = \{v_i v_{i+1} \mid 1 \leq i \leq n-1\}$$

$$E_2 = \{v_i u_i \mid 1 \leq i \leq n\}$$

$$E_3 = \{v_i w_i \mid 1 \leq i \leq n\}$$

$$E_4 = \{v_i x_i \mid 1 \leq i \leq n\}$$

Here $|G| = 4n$

$$|G| = 4n - 1$$

Define $f: V \rightarrow \{0, 1, 2, \dots, p-1\}$ by

$$f(v_i) = \{4(i-1) \mid 1 \leq i \leq n\}$$

$$f(u_i) = \{4i-3 \mid 1 \leq i \leq n\}$$

$$f(w_i) = \{4i-2 \mid 1 \leq i \leq n\}$$

$$f(x_i) = \{4i-1 \mid 1 \leq i \leq n\}$$

Then the corresponding edge labeling is as follows

$$f_{sum}^*(v_i v_{i+1}) = f(v_i) + f(v_{i+1})$$

$$= 4(i-1) + 4i$$

$$= 4i - 4 + 4i$$

$$= 8i - 4$$

$$f_{sum}^*(v_i u_i) = f(v_i) + f(u_i)$$

$$= 4(i-1) + (4i-3)$$

$$= 4i - 4 + 4i - 3$$

$$= 8i - 7$$

$$f_{sum}^*(v_i w_i) = f(v_i) + f(w_i)$$

$$= 4(i-1) + 4i - 2$$

$$= 4i - 4 + 4i - 2$$

$$= 8i - 6$$


$$f_{sum}^*(v_i x_i) = f(v_i) + f(x_i)$$

$$= 4(i - 1) + 4i - 1$$

$$= 4i - 4 + 4i - 1$$

$$= 8i - 5$$

Clearly the edge labels are distinct. Therefore pendant graph with 3 pendant is a edge sum graph.



4 Conclusions

In this paper, we proved Pendant graph with a vertex, Pendant graph with 2 vertices, Pendant graph with 3 vertices are edge sum graph. There may be interesting edge sum graphs can be constructed in future.

References

- [1] A Galian, A dyanamic survey of graph labeling, The electronic journal of combinatories (2019).
- [2] Germina K.A, Reena Sebastian, On Square Sum Graphs,Proyecciones Jouranal of Mathematics, Vol. 32 No. 2, pp. 107-117 June 2013.
- [3] R. Gopi, Odd sum labeling of Tree related graph, International Journal of Mathematics and its Application, Volume 4 Issue 4 (2016) 11-16
- [4]R. Ponraj, J. Vijaya Xavier Parthipan and R. Kala, International J. Math. Combin, Volume (4) (2010), 53-61

AUTHORS

FIRST AUTHOR – S. Alice Pappa, Associate Professor Department of Mathematics, Nazareth Margoschis College at Pillaiyanmanai, Nazareth-628617, Tamil Nadu, India. Email alicestephen8979@gmail.com

SECOND AUTHOR – R. Sweetly, Associate Professor Department of Mathematics, Nazareth Margoschis College at Pillaiyanmanai Nazareth-628617, Tamil Nadu, India. Email sweetydaniel65@gmail.com

THIRD AUTHOR - G.J. Jeba Selvi Kavitha, Department of Mathematics, Assistant Professor, St. Joseph's College of Arts and Science for Women- Hosur. Email jebaselvikavitha@gmail.com

CORRESPONDENCE AUTHOR – Jeba Selvi Kavitha G.J, Email jebaselvikavitha@gmail.com, Email vimjas1@gmail.com, CONTACT: 9486001853